arXiv:hep-ph/0202014v1 4 Feb 2002

Moduli problem and Q-ball baryogenesis in gauge-mediated SUSY breaking models

MASAHIRO KAWASAKI

Research Center for the Early Universe, School of Science, University of Tokyo
Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan

We investigate whether the Affleck-Dine mechanism can produce sufficient baryon number of the universe in the gauge-mediated SUSY breaking models, while evading the cosmological moduli problem by late-time entropy production. We find that the Q-ball formation makes the scenario difficult, irrespective of the detail mechanism of the entropy production.

PRESENTED AT

COSMO-01
Rovaniemi, Finland,
August 29 – September 4, 2001

1 Introduction

In superstring theories, there generally exist various dilaton and modulus fields. These fields (we call them "moduli") are expected to acquire masses of the order of the gravitino mass $m_{3/2}$ through some non-perturbative effects of supersymmetry (SUSY) breaking. It is well known that the moduli cause serious cosmological problem [1] because they have only gravitationally suppressed interactions with other particles and hence have long lifetimes. The moduli with mass $O(100)$ GeV decay at the Big Bang Nucleosynthesis (BBN) epoch and spoil the success of the BBN by destroying the synthesized light elements, while moduli with lighter mass ($\lesssim 1$ GeV) may overclose the universe or emit X(γ)-rays giving too many contributions to the cosmic background radiation [2].

The mass of moduli (= gravitino mass $m_{3/2}$) depends on models of SUSY breaking. In hidden sector models [3] the SUSY breaking in the hidden sector is mediated by gravitation and SUSY particles (squarks, sleptons, etc) in the observable sector as well as gravitino obtain the mass of weak scale $\sim O(100)$ GeV. On the other hand, in gauge-mediated SUSY breaking models [4], the gauge interactions mediate the SUSY breaking effects. In gauge-mediated SUSY breaking, the gravitino cannot acquires mass by the gauge interactions but only through gravitation. Thus, the mass of gravitino is much lighter ($\lesssim 1$ GeV) than that in the hidden sector models. Here we consider the moduli problem in the gauge-mediated SUSY breaking models.

In order to avoid the moduli problem, we need some huge entropy production process by which the moduli density is diluted. So far the most successful mechanism for entropy production is "thermal inflation" proposed by Lyth and Stewart [5]. The thermal inflation model in the gauge-mediated SUSY breaking models was intensively investigated in Refs. [6, 7, 8] where it was shown that the thermal inflation can solve the moduli problem. However, any process that dilutes the moduli also dilutes primordial baryon asymmetry of the universe. Thus, we must produce sufficiently large baryon asymmetry before the entropy production occurs. The promising candidate for mechanism of such efficient baryon number generation is the Affleck-Dine (AD) baryogenesis [9]. In fact, it was shown in Ref. [8] that both the present baryon asymmetry and small moduli density can be explained by the thermal inflation and the Affleck-Dine mechanism for the gauge-mediated SUSY breaking models.

However, it has been found that the dynamics of the Affleck-Dine baryogenesis is complicated by the existence of Q balls [10, 11]. In the gauge-mediated SUSY breaking models the potential for the Affleck-Dine field becomes flat at large amplitudes. For such a flat potential the Q-ball formation is inevitable [12, 13, 14]. In order to produce a large baryon number the initial amplitude of the AD field should be large, which also leads to the formation of Q balls with huge baryon(=Q) number. Since large Q balls are stable, the baryon number may be confined in the form of Q balls and there may exist very small baryon asymmetry in the cosmic plasma, which means that the baryogenesis does not work.

Unstable Q balls can provide all the charges created before the charge trapping by the produced Q balls, but rather small amplitudes of the AD field are necessary for the Q balls to decay into the ordinary baryons, nucleons. Thus, sufficient baryon number is not created from the beginning.

Therefore we study the cosmological moduli problem and baryogenesis in the gauge-mediated SUSY breaking models taking into account the Q-ball formation. It is found that the Q balls seriously affect the Affleck-Dine baryogenesis and lower its efficiency*

2 Moduli Problem

The modulus field η obtain a mass of the order of the gravitino mass $m_{3/2}$. During the primordial inflation the modulus field is expected to sit at some minimum of the effective potential determined by the Kähler potential and the Hubble parameter. In general, the minimum during the inflation deviates from the true minimum of the moduli potential at low energies and the difference of the two minimum is considered to be of the order of the gravitational scale $M(= 2.4 \times 10^{18}\text{GeV})$. After the inflation, when the Hubble parameter becomes comparable to the mass of the modulus, the modulus field begins to roll down toward the true minimum and oscillates. Then, the modulus density is estimated as

$$\frac{\rho_{mod}}{s} \simeq \frac{1}{8} T_{RH} \left(\frac{\eta_0}{M}\right)^2, \tag{1}$$

where s is the entropy density, T_{RH} is the reheating temperature and η_0 is the initial amplitude of the modulus oscillation ($\eta_0 \sim M$). In deriving Eq.(1), we have assumed that the modulus mass is equal to $m_{3/2}$ and the reheating takes after the moduli field starts the oscillation. Since T_{RH} should be higher than about 10 MeV to keep the success of the BBN, the moduli to entropy ratio is bounded from below,

$$\frac{\rho_{mod}}{s} \gtrsim 1.25 \times 10^{-3}\text{GeV}. \tag{2}$$

The decay rate of the modulus is very small because it has only gravitationally suppressed interaction. The lifetime is roughly estimated as

$$\tau_\eta \sim 10^{18}\,\text{sec} \left(\frac{m_{3/2}}{100\text{MeV}}\right)^{-3}. \tag{3}$$

Thus, for $m_{3/2} \lesssim 100$ MeV, the lifetime is longer than the age of the universe and its present density much larger than the critical density which is given by $\rho_c/s_0 = 3.6 \times 10^{-9} h^2\text{GeV}$, where h is the present Hubble parameter in units of 100km/sec/Mpc and $s_0(\simeq 2.8 \times 10^3\ \text{cm}^{-3})$ is the present entropy density. The modulus with larger mass (100 MeV $\lesssim m_{3/2} \lesssim$ 1 GeV) decays into photons whose flux exceeds the observed background X(or γ)-rays. Therefore the modulus is cosmological disaster and should be diluted by some entropy production process.

---

*The detailed analysis is found in Ref. [15].

3 Affleck-Dine mechanism and Q-ball formation

In the Minimal Supersymmetric Standard Model (MSSM), there exist flat directions, along which there are no classical potentials. Since flat directions consist of squarks and/or sleptons, they carry baryon and/or lepton numbers, and can be identified as the Affleck-Dine field. These flat directions are lifted by SUSY breaking effects. In the gauge-mediated SUSY breaking models, the potential of a flat direction is parabolic at the origin, and almost flat beyond the messenger scale:

$$V_{gauge} \sim \left\{ \begin{array}{ll} m_\phi^2 |\Phi|^2 & \Phi \ll M_S \\ M_F^4 \log \frac{|\Phi|^2}{M_S^2} & \Phi \gg M_S \end{array} \right. , \tag{4}$$

where M_S is the messenger mass scale.

Since the gravity always exists, flat directions are also lifted by the gravity-mediated SUSY breaking effects:

$$V_{grav} = m_{3/2}^2 \left[1 + K \log \left(\frac{|\Phi|^2}{M} \right) \right] |\Phi|^2, \tag{5}$$

where K is the numerical coefficient of the one-loop corrections. This term can be dominant only at high energy scales because of the small gravitino mass $\lesssim O(1\text{GeV})$.

In addition, there is a thermal effect on the potential, which appears at two-loop order, as pointed out in Ref. [16]. This comes from the fact that the running of the gauge coupling $g(T)$ is modified by integrating out heavy particles which directly couples with the AD field. This contribution is given by

$$V_T \sim T^4 \log \frac{|\Phi|^2}{T^2}. \tag{6}$$

The baryon number is usually created just after the AD field starts coherent rotation in the potential, and its number density n_B is estimated as

$$n_B(t_{osc}) \simeq \varepsilon \omega \phi_{osc}^2, \tag{7}$$

where $\varepsilon(\lesssim 1)$ is the ellipticity parameter, which represents the strongness of the A-term, and ω and ϕ_{osc} are the angular velocity and the amplitude of the AD field at the beginning of the oscillation (rotation) in its effective potential.

Actually, however, the AD field feels spatial instabilities during its coherent oscillation, and deforms into nontopological solitons, Q balls [10, 11, 12]. From numerical calculations [12, 14], Q balls absorb almost all the baryon charges which the AD field obtains, and the typical charge is estimated as [14]

$$Q \simeq \beta \left(\frac{\phi_{osc}}{M_F} \right)^4, \tag{8}$$

where $\beta \approx 6 \times 10^{-4}$. Consequently, the present baryon asymmetry should be explained by the charges which come out of the Q balls through the evaporation, diffusion, and decay of Q balls.

In the case of the unstable Q balls, they decay into nucleons and light scalar particles. Since the temperature at the BBN time is very low (~ 1 MeV), Q balls cannot decay into light scalars. This rate is thus given by [17]

$$\frac{dQ}{dt} \lesssim \frac{\omega^3 A}{192\pi^2}, \tag{9}$$

where A is a surface area of the Q ball.

In the case of the stable Q balls, the evaporation is the only way to extract the baryon charges from Q balls. The total evaporated charge from the Q ball is estimated as [18, 19, 14],

$$\Delta Q \sim 10^{15} \left(\frac{m_\phi}{\text{TeV}}\right)^{-2/3} \left(\frac{M_F}{10^6\text{GeV}}\right)^{-1/3} Q^{1/12}. \tag{10}$$

Hence the baryon number density is suppressed by the factor $\Delta Q/Q$, in comparison with the case of no stable Q-ball production.

On the other hand, where V_{grav} dominates the potential at larger scales, the gravity-mediation type Q balls ('new' type) are produced [13], if K is negative, while, if K is positive, it is not until the AD field enters V_{gauge} dominant region that it feels instabilities, and the gauge-mediation type Q balls are produced (the delayed Q balls) [14]. Notice that the sign of K is in general indefinite in the gauge-mediated SUSY breaking models.

When the AD field starts to oscillate in the V_{grav}-dominant region, where $H_{osc} \sim \omega \sim m_{3/2}$, the baryon number is produced as $n_B \simeq \varepsilon m_{3/2} \phi_{osc}$. For the negative K case, the 'new-type' Q balls are created, and its charge is written as

$$Q \simeq \tilde{\beta} \left(\frac{\phi_{osc}}{m_{3/2}}\right)^2, \tag{11}$$

where $\tilde{\beta} \simeq 6 \times 10^{-3}$. This type of the Q ball is also stable against the decay into nucleons, and the amount of the baryons in the present universe is explained by the charge evaporation from the Q balls. The charge evaporated from the Q ball is estimated as [13]

$$\Delta Q \sim 2.2 \times 10^{20} \left(\frac{m_{3/2}}{100\text{keV}}\right)^{-1/3} \left(\frac{m_\phi}{\text{TeV}}\right)^{-2/3}. \tag{12}$$

4 Baryogenesis and the moduli problem

As we mentioned in the Introduction, the late-time entropy production necessary for the dilution of the moduli also dilutes the baryon numbers created earlier very seriously, but the sufficient numbers could remain, if the Q-ball production is not taken into account. We will see that the Q-ball formation put very serious restriction on the efficiency of the AD baryogenesis, whether the produced Q balls are stable or not.

4.1 Stable Q balls

Since the baryon number is supplied only by the evaporation from stable Q balls, it should be suppressed by the factor $\Delta Q/Q$, compared with no Q-ball formation. We will show that this fact considerably reduces the power of the AD baryogenesis.

4.1.1 Gauge-mediation type Q balls when the zero-temperature potential is dominated

If the V_{gauge} dominates over the thermal logarithmic potential V_T, $M_F \gtrsim T_{osc}$, so that the initial amplitude is constrained as $\phi_{osc} \gtrsim (T_{RH}/M_F)^2 M$. V_{gauge} also dominates over V_{grav}, which leads to the condition $\phi_{osc} \lesssim M_F^2/m_{3/2}$. Combining these two equations, we have $T_{RH} \lesssim M_F^2/\sqrt{m_{3/2}M}$.

From the stability ($M_Q/Q < 1$ GeV) and survival condition ($\Delta Q < Q$), the maximal baryon-to-entropy ratio is

$$Y_B \quad \lesssim \quad 6.2 \times 10^{-27} \varepsilon \left(\frac{\Omega_{mod} h^2}{0.2}\right) \left(\frac{m_\phi}{\text{TeV}}\right)^{-2/3}. \tag{13}$$

Therefore, the baryon-to-entropy ratio is too small to explain the present value of the order 10^{-10}.

4.1.2 Gauge-mediation type Q balls when the thermal logarithmic potential is dominated

In this case, V_T is dominant over V_{gauge}, and the AD field starts to oscillate when $H_{osc} \sim T_{osc}^2/\phi_{osc}$. We should also use the charge of the formed Q ball as $Q \simeq \beta(\phi_{osc}/T_{osc})^4$. Therefore, the fraction of the evaporated charge is written as

$$\frac{\Delta Q}{Q} \quad \sim \quad 8.9 \times 10^{17} \left(\frac{m_\phi}{\text{TeV}}\right)^{-2/3} \left(\frac{M_F}{10^6\text{GeV}}\right)^{-1/3} \left(\frac{T_{RH}}{M}\right)^{11/3} \left(\frac{\phi_{osc}}{M}\right)^{-11/2}. \tag{14}$$

Then the baryon-to-entropy ratio becomes

$$Y_B \quad \simeq \quad 5.3 \times 10^{-10} \varepsilon \left(\frac{\Omega_{mod} h^2}{0.2}\right) \left(\frac{m_\phi}{\text{TeV}}\right)^{-2/3} \left(\frac{M_F}{10^6\text{GeV}}\right)^{-1/3} \left(\frac{T_{RH}}{M}\right)^{5/3} \left(\frac{\phi_{osc}}{M}\right)^{-3/2} \tag{15}$$

We plot the baryon-to-entropy ratio in the function of $m_{3/2}$ in Fig. 1. We can thus marginally explain the present value ($Y_B \gtrsim 10^{-11}$) for $m_{3/2} \lesssim 100$ keV, but the reheating temperature is $\sim 2.0 \times 10^{13}$ GeV, so high for natural inflation models to provide. It is noticed that Y_B is maximized at $\Delta Q \sim Q$.



Figure 1: Largest possible baryon-to-entropy ratio in the stable Q-ball scenario in the thermal logarithmic potential.

4.1.3 Delayed Q balls when the zero-temperature potential is dominated

Both the AD field and moduli fields start to oscillate when $H_{osc} \sim m_{3/2}$, the ratio of the baryon number and the energy density of the moduli stays constant to the present, the baryon number is given by $n_B \simeq \varepsilon m_{3/2}\phi_{osc}^2 \Delta Q/Q$, and the baryon-to-entropy ratio becomes

$$Y_B \sim 2.8 \times 10^{-24}\varepsilon \left(\frac{\Omega_{mod}h^2}{0.2}\right)\left(\frac{m_\phi}{\mathrm{TeV}}\right)^{-2/3}\left(\frac{m_{3/2}}{100\mathrm{keV}}\right)^{8/3}\left(\frac{M_F}{10^6\mathrm{GeV}}\right)^{-4}\left(\frac{\phi_{osc}}{M}\right)^2. \quad (16)$$

Figure 2 shows the maximum value of the baryon-to-entropy ratio. As can be seen, this scenario is marginally successful ($Y_B \sim 10^{-11}$) only for $m_{3/2} \sim 200$ keV. Notice that, the value of Y_B is the same as that in the thermal logarithmic potential, to be considered in the next subsection, for $m_{3/2} \lesssim 0.16$ GeV.

4.1.4 Delayed Q balls when the thermal logarithmic potential is dominated

In the case that V_T is dominant over V_{gauge}, the maximum value of the baryon-to-entropy ratio is also shown in Fig. 2. We can see that Y_B is marginally enough ($Y_B \sim 10^{-11}$) only for $m_{3/2} \simeq 200$ keV. Notice that the largest possible value of Y_B is achieved when $\Delta Q \sim Q$.



Figure 2: Largest possible baryon-to-entropy ratio in the stable delayed Q-ball scenario. Notice that it can be applied in the zero-temperature logarithmic potential V_{gauge} for $m_{3/2} \lesssim 0.16$ GeV, while all ranges of $m_{3/2}$ can be applied in the thermal logarithmic potential V_T.

4.1.5 New type Q balls

From Eqs.(11) and (12), we have

$$\frac{\Delta Q}{Q} \simeq 6.1 \times 10^{-23} \left(\frac{m_\phi}{\text{TeV}}\right)^{-2/3} \left(\frac{m_{3/2}}{100\text{keV}}\right)^{5/3} \left(\frac{\phi_{osc}}{M}\right)^{-2}. \tag{17}$$

This leads to the baryon-to-entropy ratio as

$$Y_B \lesssim 8.8 \times 10^{-28} \varepsilon \left(\frac{\Omega_{mod} h^2}{0.2}\right) \left(\frac{m_\phi}{\text{TeV}}\right)^{-2/3} \left(\frac{m_{3/2}}{100\text{keV}}\right)^{2/3}, \tag{18}$$

which is too small to explain the present value.

4.2 Unstable Q balls

In the case of the unstable Q balls which decay into nucleons, it may destroy light elements synthesized at the BBN. Thus, a new constraint which the Q ball should decay before the BBN (∼ 1 sec), must be imposed.

4.2.1 Gauge-mediation type Q balls when the zero-temperature potential is dominated

There are several condition to be imposed. The first is the condition that the Q ball is unstable, given by

$$\frac{\phi_{osc}}{M} \lesssim 2.6 \times 10^{-6} \left(\frac{M_F}{10^6 \mathrm{GeV}}\right)^2 . \tag{19}$$

Second, the decay of the Q ball must be completed until the BBN, otherwise it would spoil the success of the BBN, so that the life time τ_Q should be less than about 1 sec. Thus, we have the following constraint:

$$\frac{\phi_{osc}}{M} \lesssim 1.0 \times 10^{-6} \left(\frac{M_F}{10^6 \mathrm{GeV}}\right)^{6/5} . \tag{20}$$

The largest possible value of Y_B is obtained by the conditions Eq.(20) and $M_F \lesssim (m_{3/2} M)^{1/2}$, and will be written as

$$Y_B \lesssim 2.8 \times 10^{-19} \varepsilon \left(\frac{\Omega_{mod} h^2}{0.2}\right) \left(\frac{m_{3/2}}{100 \mathrm{keV}}\right)^{4/5} . \tag{21}$$

This is thus too small to explain the present value $\sim 10^{-10}$.

4.2.2 Gauge-mediation type Q balls when the thermal logarithmic potential is dominated

Since the Q-ball charge is expressed as $Q \simeq \beta(\phi_{osc}/M)^6 (T_{RH}/M)^{-4}$, the unstable condition, $M_F Q^{-1/4} \gtrsim 1$ GeV, is given by

$$\frac{\phi_{osc}}{M} \lesssim 3.4 \times 10^4 \left(\frac{M_F}{10^6 \mathrm{GeV}}\right)^{2/3} \left(\frac{T_{RH}}{M}\right)^{2/3} , \tag{22}$$

while the lifetime condition that the Q ball decays before the BBN time ($\sim$ 1 sec), is written as

$$\frac{\phi_{osc}}{M} \lesssim 4.0 \times 10^3 \left(\frac{M_F}{10^6 \mathrm{GeV}}\right)^{2/15} \left(\frac{T_{RH}}{M}\right)^{2/3} . \tag{23}$$

From the above conditions, we have the baryon-to-entropy ratio as

$$Y_B \lesssim 8.7 \times 10^{-11} \varepsilon \left(\frac{\Omega_{mod} h^2}{0.2}\right) \left(\frac{m_{3/2}}{100 \mathrm{keV}}\right)^{-2/5} . \tag{24}$$

We plot the largest possible value of Y_B [Eq.(24)] in the function of $m_{3/2}$ in Fig. 3. As can be seen, we can explain the present value in the range $m_{3/2} = 10 - 100$ keV. However, the reheating temperature should be as high as 5.6×10^{12} GeV, which may be rather higher for the actual inflation models.



Figure 3: Largest possible baryon-to-entropy ratio in the unstable Q-ball scenario in the thermal logarithmic potential.

4.2.3 Delayed Q balls

Firstly, it is found that the unstable condition can be expressed as $m_{3/2} \gtrsim 0.16$ GeV both when $V_T \lesssim V_{gauge}$ and when $V_T \gtrsim V_{gauge}$. Therefore, in either case, we can estimate the baryon-to-entropy ratio as

$$Y_B \lesssim 4.5 \times 10^{-17} \varepsilon \left(\frac{\Omega_{mod} h^2}{10^{-9}} \right) \left(\frac{m_{3/2}}{0.16 \text{GeV}} \right)^{-1} \left(\frac{\phi_{osc}}{M} \right)^2, \tag{25}$$

where $\Omega_{mod} h^2 \lesssim 10^{-9}$ for $m_{3/2} \gtrsim 0.16$ GeV from the X(γ)-ray constraint. This is too small to explain the present value, even if $\phi_{osc} \sim M$.

5 Conclusion

We have investigate the possibility of the AD baryogenesis in the gauge-mediated SUSY breaking scenario, while evading the cosmological moduli problem by the late-time entropy production. In all the cases, the Q-ball formation makes the efficiency of the baryon number production considerably be diminish. In the zero-temperature potential V_{gauge}, whether the produced Q balls are stable or not, the largest possible baryon-to-entropy ratio is too small to explain the present value. This completely kills the successful situations considered in Ref. [8].

We have also found that there are some marginally successful situations when we take into account of the thermal effects on the effective potential of the AD field. However, these successful situations require very high reheating temperatures such as $10^{12} - 10^{16}$ GeV, which may be impossible to achieve in the actual inflation models.

In the delayed Q-ball formation case, we have found that enough baryon-to-entropy ratio can be created, even in the zero-temperature potential V_{gauge} is dominant over the thermal logarithmic potential V_T. It might be the unique solution for the AD baryogenesis with solving the cosmological moduli problem, although the scale of M_F is rather low.

In addition, successful situations above need the following conditions: the large initial amplitude such as $\phi_{osc} \simeq M$, and $\varepsilon \simeq 1$. This is realized if the A-terms, which make the AD field rotate in the effective potential, originate from some Kähler potential. Then, $\varepsilon \sim (\phi/M)^{\gamma} \sim 1$ for $\phi \sim M$, where $\gamma > 2$.

ACKNOWLEDGEMENTS

This work is supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture of Japan, Priority Area "Supersymmetry and Unified Theory of Elementary Particles" (No. 707).

References

[1] G.D. Coughlan, W. Fischler, E.W. Kolb, S.Raby and G.G. Ross, Phys. Lett. **131B**, 59 (1983).

[2] M. Kawasaki and T. Yanagida, Phys. Lett. **B399**, 45 (1997).

[3] For a review, see, e.g. H.P. Nilles, Phys. Rep. **110**, 1 (1984).

[4] For a review, see, e.g. G.F. Giudice and R. Rattazzi, Phys. Rep. **322**, 419 (1999).

[5] D.H. Lyth and E.D. Stewart, Phys. Lett. **75**, 201 (1995);
Phys. Rev. **D53**, 1784 (1996).

[6] J. Hashiba, M. Kawasaki and T. Yanagida, Phys. Rev. Lett. **79**, 4525 (1997).

[7] T. Asaka, J. Hashiba, M. Kawasaki and T. Yanagida, Phys. Rev. **D58**, 083509 (1998).

[8] T. Asaka and M. Kawasaki, Phys. Rev. **D60**, 123509 (1999).

[9] I. Affleck and M. Dine, Nucl. Phys. **B249**, 361 (1985).

[10] A. Kusenko and M. Shaposhnikov, Phys. Lett. **B418**, 46 (1998).

[11] K. Enqvist and J. McDonald, Nucl. Phys. **B538**, 321 (1999).

[12] S. Kasuya and M. Kawasaki, Phys. Rev. **D61**, 041301 (2000).

[13] S. Kasuya and M. Kawasaki, Phys. Rev. Lett. **85**, 2677 (2000).

[14] S. Kasuya and M. Kawasaki, to appear in Phys. Rev. **D**, hep-ph/0106119.

[15] S. Kasuya, M. Kawasaki and F. Takahashi, hep-ph/0108171, to appear in Phys. Rev. **D**.

[16] A. Anisimov and M. Dine, hep-th/0008058.

[17] A. Cohen, S. Coleman, H. Georgei, and A. Manohar, Nucl. Phys. **B272**, 301 (1986).

[18] M. Laine, and M. Shaposhnikov, Nucl. Phys. **B532**,376 (1998).

[19] R. Banerjee and K. Jedamzik, Phys. Lett. **B484**, 278 (2000).